



**DIGITAL ACCESS TO
SCHOLARSHIP AT HARVARD**
DASH.HARVARD.EDU



HARVARD LIBRARY
Office for Scholarly Communication

Author Correction: Multiplexed imaging of high-density libraries of RNAs with MERFISH and expansion microscopy

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Wang, Guiping, Jeffrey R. Moffitt, and Xiaowei Zhuang. 2018. "Author Correction: Multiplexed imaging of high-density libraries of RNAs with MERFISH and expansion microscopy." Scientific Reports 8 (1): 6487. doi:10.1038/s41598-018-24844-8. http://dx.doi.org/10.1038/s41598-018-24844-8 .
Published Version	doi:10.1038/s41598-018-24844-8
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:37068317
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

SCIENTIFIC REPORTS

OPEN

Author Correction: Multiplexed imaging of high-density libraries of RNAs with MERFISH and expansion microscopy

Guiping Wang, Jeffrey R. Moffitt & Xiaowei Zhuang 

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-018-22297-7>, published online 19 March 2018

This Article contains errors in the Methods section under subheading ‘Encoding probe staining’.

“Then 30 μ L of $\sim 300 \mu$ M encoding probes and 3.3 μ M of poly(dT) LNA anchor probe (a 20-nt sequence of alternating dT and thymidine-locked nucleic acid (dT+) with a 20-nt reverse complement of a readout sequence and a 5'-acrydite modification (Integrated DNA Technologies)) in encoding hybridization buffer was added to the surface of Parafilm (Bemis) and was covered with a cell-containing coverslip.”

should read:

“Then 30 μ L of $\sim 50 \mu$ M encoding probes and 3.3 μ M of poly(dT) LNA anchor probe (a 20-nt sequence of alternating dT and thymidine-locked nucleic acid (dT+) with a 20-nt reverse complement of a readout sequence and a 5'-acrydite modification (Integrated DNA Technologies)) in encoding hybridization buffer was added to the surface of Parafilm (Bemis) and was covered with a cell-containing coverslip.”



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018

Howard Hughes Medical Institute, Department of Chemistry and Chemical Biology, Department of Physics, Harvard University, Cambridge, MA, 02138, USA. Correspondence and requests for materials should be addressed to X.Z. (email: zhuang@chemistry.harvard.edu)